

FIRST EXTENSION GROUPS OF VERMA MODULES AND R -POLYNOMIALS

NORIYUKI ABE

ABSTRACT. We study the first extension groups between Verma modules. There was a conjecture which claims that the dimensions of the higher extension groups between Verma modules are the coefficients of R -polynomials defined by Kazhdan-Lusztig. This conjecture was known as the Gabber-Joseph conjecture (although Gebber and Joseph did not state.) However, Boe gives a counterexample to this conjecture. In this paper, we study how far are the dimensions of extension groups from the coefficients of R -polynomials.

1. INTRODUCTION

The category $\mathcal{O}$ is introduced by Bernstein-Gelfand-Gelfand and plays an important role in the representation theory. One of the most important objects in $\mathcal{O}$ are the Verma modules and they are deeply investigated.

In this paper, we consider the Ext^i -groups between Verma modules. If $i = 0$, Verma [Ver68] and Bernstein-Gelfand-Gelfand [BGG71] determines the dimension of this group. There are many studies about higher extension groups. One of these studies is a work of Gebber-Joseph. They proved some inequality between the dimensions of extension groups. After that, it is conjectured that this inequality is, in fact, an equality. Although not actually stated in [GJ81], this conjecture is known as the Gebber-Joseph conjecture. If this conjecture is true, then the dimension of extension groups are the coefficients of R -polynomial. However, Boe gives a counterexample to this conjecture [Boe92].

This conjecture is false even in the case of $i = 1$. In this paper, we consider how far the dimensions of extension groups from the coefficients of R -polynomials. Mazorchuk gives a formula of the dimension of the first extension group between Verma modules in a special case [Maz07, Theorem 32]. Our formula is a generalization of his formula.

Now we state our main theorem. Let $\mathfrak{g}$ be a semisimple Lie algebra over an algebraic closed field K of characteristic zero. Fix its Borel subalgebra $\mathfrak{b}$ and a Cartan subalgebra $\mathfrak{h}$. Let Δ be the root system and ρ the half sum of positive roots. Fix a dominant integral element $\lambda \in \mathfrak{h}^*$. (It is sufficient to consider the integral case by a result of Soergel [Soe90, Theorem 11].) Let $M(x\lambda)$ be the Verma module with highest weight $x\lambda - \rho$ for $x \in W$. Then by a result of Verma [Ver68] and Bernstein-Gelfand-Gelfand [BGG71], if $x \geq y$, then there exists the unique (up to nonzero constant multiple) injective homomorphism $M(x\lambda) \rightarrow M(y\lambda)$. Hence we can regard $M(x\lambda)$ as a submodule of $M(y\lambda)$. Then $M(w_0\lambda)$ is a submodule of $M(x\lambda)$ for all $x \in W$ where w_0 is the longest Weyl element. Hence we have the homomorphism $\text{Ext}^1(M(x\lambda), M(y\lambda)) \rightarrow \text{Ext}^1(M(w_0\lambda), M(y\lambda))$. Let $V_\lambda(x, y)$ be

the image of this homomorphism. We denote the unit element of W by e . Put $S_\lambda = \{s \in S \mid s(\lambda) = \lambda\}$. For integral $\lambda, \mu \in \mathfrak{h}^*$, let T_λ^μ be the translation functor from λ to μ .

Theorem 1.1 (Proposition 3.14, Theorem 4.3, Theorem 5.1). *Let λ be a dominant integral element.*

- (1) *If λ is regular, then $V_\lambda(w_0, e)$ has a structure of W -module and it is isomorphic to $\mathfrak{h}^*$. For $s \in S$, we denote the element in $V_\lambda(w_0, e)$ corresponding to the simple root whose reflection is s by $v_s \in V_\lambda(x, y)$.*
- (2) *Assume that λ is regular. For $x, y \in W$ and $s \in S$ such that $xs > x$, we have the following formula.*
 - (a) *If $ys < y$, then $V_\lambda(xs, y) = s(V_\lambda(x, ys))$.*
 - (b) *If $ys > y$, then $V_\lambda(xs, y) = Kv_s + s(V_\lambda(x, y))$.*
- (3) *For general λ , the translation functor T_ρ^λ induces a linear map $V_\rho(w_0, e) \rightarrow V_\lambda(w_0, e)$. The kernel of this linear map is $\sum_{s \in S_\lambda} Kv_s$ and $T_\rho^\lambda(V_\rho(x, y)) = V_\lambda(x, y)$.*

Since $V(x, x) = 0$ for all $x \in W$, we can determine the space $V_\lambda(x, y)$ inductively. In particular, we can calculate the dimension of $V_\lambda(x, y)$. Namely, we can get the following theorem. For $x, y \in W$, let $R_{x, y}$ be the polynomial defined in [KL79, §2]. Let $\ell(x)$ be the length of $x \in W$.

Theorem 1.2 (Theorem 4.4). *Assume that λ is regular. Then $\dim V_\lambda(x, y)$ is the coefficient of q in $(-1)^{\ell(y) - \ell(x) - 1} R_{y, x}(q)$.*

In other words, $V_\lambda(x, y)$ satisfies the Gebber-Joseph conjecture. Notice that the homomorphism $\text{Ext}^1(M(x\lambda), M(y\lambda)) \rightarrow \text{Ext}^1(M(w_0\lambda), M(\lambda))$ is not injective. It is easy to see that the kernel is isomorphic to $\text{Hom}(M(x\lambda), M(\lambda)/M(y\lambda))$ if $x \geq y$ (Lemma 2.1).

We summarize the contents of this paper. In Section 2, we gather preliminaries and prove some easy facts. In Section 3, we define a W -module structure on $\text{Ext}^1(M(w_0\lambda), M(\lambda))$ for a dominant integral regular λ . We also prove this module is isomorphic to $\mathfrak{h}^*$. The proof of the main theorem in the regular case is done in Section 4. We finish the proof of the main theorem in Section 5.

2. PRELIMINARIES

Let $\mathfrak{g}$ be a semisimple Lie algebra over an algebraic closed field K of characteristic zero. Fix a Borel subalgebra $\mathfrak{b}$ and a Cartan subalgebra $\mathfrak{h} \subset \mathfrak{b}$. These determine the BGG category $\mathcal{O}$ [BGG76]. We denote $\text{Hom}_{\mathcal{O}}$ (resp. $\text{Ext}_{\mathcal{O}}^i$) by Hom (resp. Ext^i). Denote the Weyl group of $\mathfrak{g}$ by W and let S be the set of its simple reflections. For $w \in W$, let $\ell(w)$ be its length. For $\lambda \in \mathfrak{h}^*$, let $\mathcal{O}_\lambda$ be the full-subcategory of $\mathcal{O}$ consisting of objects which have the generalized infinitesimal character λ .

In the rest of this paper, we only consider the objects which has a integral generalized infinitesimal character. By [Soe90, Theorem 11], it is sufficient to consider only the integral case.

For $\lambda \in \mathfrak{h}^*$, let $M(\lambda)$ be the Verma module with highest weight $\lambda - \rho$ where ρ is the half sum of positive roots. Assume that λ is dominant integral. Set $S_\lambda = S \cap \text{Stab}_W \lambda$. Then $W_{S_\lambda} = \text{Stab}_W \lambda$ is generated by S_λ . Put $W(S_\lambda) = \{w \in W \mid ws > w \text{ for all } s \in S_\lambda\}$. This gives a complete representative set of W/W_{S_λ} . For $x, y \in W(S_\lambda)$, $\text{Hom}(M(x\lambda), M(y\lambda)) \neq 0$ if and only if $x \geq y$. Moreover, if $x \geq$

y , then $\text{Hom}(M(x\lambda), M(y\lambda))$ is one-dimensional and the nonzero homomorphisms from $M(x\lambda)$ to $M(y\lambda)$ are injective. Hence we can regard $M(x\lambda)$ as a submodule of $M(y\lambda)$. In the rest of this paper, we regard $M(x\lambda)$ as a submodule of $M(\lambda)$. Then $M(w_0\lambda)$ is a submodule of $M(x\lambda)$ for all $x \in W$.

Let $\lambda, \mu \in \mathfrak{h}^*$ be integral dominant elements. Then the translation functor $T_\lambda^\mu: \mathcal{O}_\lambda \rightarrow \mathcal{O}_\mu$ is defined [Jan79].

Assume that λ is a regular integral dominant element. Then for $s \in S$, the wall-crossing functor θ_s^λ is defined by $\theta_s^\lambda = T_\mu^\lambda T_\lambda^\mu$ where $\text{Stab}_W \mu = \{e, s\}$. It is known that this functor is independent of μ . By the definition of θ_s^λ , there are natural transformations $\text{Id} \rightarrow \theta_s^\lambda$ and $\theta_s^\lambda \rightarrow \text{Id}$. Put $C_s^\lambda = \text{Cok}(\text{Id} \rightarrow \theta_s^\lambda)$ and $K_s^\lambda = \text{Ker}(\theta_s^\lambda \rightarrow \text{Id})$. Then C_s^λ (resp. K_s^λ) gives a right (resp. left) exact functor from $\mathcal{O}_\lambda$ to $\mathcal{O}_\lambda$. By the self-adjointness of θ_s^λ , $(C_s^\lambda, K_s^\lambda)$ is an adjoint pair. Moreover, the derived functor LC_s gives an auto-equivalence of the derived category $D^b(\mathcal{O}_\lambda)$ and its quasi-inverse is RK_s . For the action of C_s, K_s on Verma modules, the following formulas hold: Let $x \in W$ and $s \in S$ such that $xs > x$. Then $C_s(M(x)) = M(xs)$, $K_s(M(xs)) = M(x)$ and there exists an exact sequence $0 \rightarrow M(x)/M(xs) \rightarrow C_s(M(xs)) \rightarrow M(xs) \rightarrow 0$. Moreover, we have $L^k C_s(M(x)) = L^k C_s(M(xs)) = R^k K_s(M(xs)) = 0$ for all $k \geq 1$.

Let λ be the dominant integral element of $\mathfrak{h}^*$. Set

$$V_\lambda(x, y) = \text{Im}(\text{Ext}^1(M(x\lambda), M(y\lambda)) \rightarrow \text{Ext}^1(M(w_0\lambda), M(\lambda)))$$

for $x, y \in W$. Put

$$V_\lambda = V_\lambda(w_0, e).$$

(1) of the following lemma is a part of the argument of the proof of [Maz07, Lemma 33]. (2) and (3) follows from (1).

Lemma 2.1. *Let λ be a dominant integral element.*

- (1) *Let P be the projective cover of $M(w_0\lambda)$. Let $M_i \subset P$ be the filtration such that $M_i/M_{i-1} \simeq \bigoplus_{\ell(w)=i, w \in W(S_\lambda)} M(w\lambda_0)$. Then we have*

$$\begin{aligned} \text{Ext}^1(M(x\lambda), M(\lambda)) &\xleftarrow{\sim} \text{Hom}(M(x\lambda), P/M_0) \\ &\xleftarrow{\sim} \text{Hom}(M(x\lambda), M_1/M_0) \simeq \bigoplus_{s \in S \setminus S_\lambda} \text{Hom}(M(x\lambda), M(s\lambda)). \end{aligned}$$

- (2) *We have $\dim V_\lambda = \#(S \setminus S_\lambda)$.*

- (3) *For $x, y \in W$ such that $x \geq y$, the kernel of $\text{Ext}^1(M(x\lambda), M(y\lambda)) \rightarrow \text{Ext}^1(M(w_0\lambda), M(\lambda))$ is isomorphic to $\text{Hom}(M(x\lambda), M(\lambda)/M(y\lambda))$.*

Proof. (1) follows from the proof of [Maz07, Lemma 33]. (2) follows from (1). We prove (3). We have the long exact sequence

$$\begin{aligned} 0 \rightarrow \text{Hom}(M(x\lambda), M(y\lambda)) &\rightarrow \text{Hom}(M(x\lambda), M(\lambda)) \\ \rightarrow \text{Hom}(M(x\lambda), M(\lambda)/M(y\lambda)) &\rightarrow \text{Ext}^1(M(x\lambda), M(y\lambda)) \rightarrow \text{Ext}^1(M(x\lambda), M(\lambda)) \end{aligned}$$

The morphism $\text{Hom}(M(x\lambda), M(y\lambda)) \rightarrow \text{Hom}(M(x\lambda), M(\lambda))$ is isomorphic by the classification of homomorphism between Verma modules.

Hence it is sufficient to prove that $\text{Ext}^1(M(x\lambda), M(\lambda)) \rightarrow \text{Ext}^1(M(w_0\lambda), M(\lambda))$ is injective. This follows from (1). $\square$

3. WEYL GROUP ACTION

In this section, fix a dominant regular integral element λ . Put $V = V_\lambda$, $V(x, y) = V_\lambda(x, y)$, $\theta_s = \theta_s^\lambda$, $C_s = C_s^\lambda$ and $M(x) = M(x\lambda)$. Since $M(x) \rightarrow \theta_s(M(x)) \rightarrow M(x)$ is zero, we have $C_s(M(x)) \rightarrow M(x)$.

We begin with the following lemma.

Lemma 3.1. *For M_1, M_2, N_1, N_2 , take $a_i \in \text{Ext}^1(M_i, N_i)$ for $i = 1, 2$. Take an exact sequence $0 \rightarrow N_i \xrightarrow{k_i} X_i \xrightarrow{p_i} M_i \rightarrow 0$ corresponding to a_i . Then for $f: N_1 \rightarrow N_2$ and $g: M_1 \rightarrow M_2$, there exists φ such that the following diagram commutes if and only if $f_*a_1 = g^*a_2$:*

$$\begin{array}{ccccccccc} 0 & \longrightarrow & N_1 & \longrightarrow & X_1 & \longrightarrow & M_1 & \longrightarrow & 0 \\ & & \downarrow f & & \downarrow \varphi & & \downarrow g & & \\ 0 & \longrightarrow & N_2 & \longrightarrow & X_2 & \longrightarrow & M_2 & \longrightarrow & 0. \end{array}$$

Proof. First assume that there exists φ . Consider the following digram:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & N_1 & \longrightarrow & X_1 & \longrightarrow & M_1 & \longrightarrow & 0 \\ & & \downarrow f & & \downarrow & & \parallel & & \\ 0 & \longrightarrow & N_2 & \longrightarrow & X' & \longrightarrow & M_1 & \longrightarrow & 0, \end{array}$$

where the left square is a push-forward. Then the exact sequence $0 \rightarrow N_2 \rightarrow X' \rightarrow M_1 \rightarrow 0$ corresponds to f_*a_1 and the existence of φ implies the existence of φ' such that the following diagram commutes:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & N_2 & \longrightarrow & X' & \longrightarrow & M_1 & \longrightarrow & 0 \\ & & \parallel & & \downarrow \varphi' & & \downarrow g & & \\ 0 & \longrightarrow & N_2 & \longrightarrow & X_2 & \longrightarrow & M_2 & \longrightarrow & 0. \end{array}$$

The same argument implies the existence of the following digram:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & N_2 & \longrightarrow & X' & \longrightarrow & M_1 & \longrightarrow & 0 \\ & & \parallel & & \downarrow \varphi'' & & \parallel & & \\ 0 & \longrightarrow & N_2 & \longrightarrow & X'' & \longrightarrow & M_1 & \longrightarrow & 0, \end{array}$$

where $0 \rightarrow N_2 \rightarrow X'' \rightarrow M_1 \rightarrow 0$ corresponds to g^*a_2 . The morphism φ'' should be isomorphism by 5-Lemma. Hence we have $f_*a_1 = g^*a_2$.

Conversely, we assume that $f_*a_1 = g^*a_2$. Consider the following diagram:

$$\begin{array}{ccccccc}
\mathrm{Hom}(M_1, N_2) & \xrightarrow{k_{2*}} & \mathrm{Hom}(M_1, X_2) & \xrightarrow{p_{2*}} & \mathrm{Hom}(M_1, M_2) & \xrightarrow{\delta_2} & \mathrm{Ext}^1(M_1, N_2) \\
\downarrow p_1^* & & \downarrow p_1^* & & \downarrow p_1^* & & \downarrow p_1^* \\
\mathrm{Hom}(X_1, N_2) & \xrightarrow{k_{2*}} & \mathrm{Hom}(X_1, X_2) & \xrightarrow{p_{2*}} & \mathrm{Hom}(X_1, M_2) & \xrightarrow{\delta_1} & \mathrm{Ext}^1(X_1, N_2) \\
\downarrow k_2^* & & \downarrow k_2^* & & \downarrow k_2^* & & \\
\mathrm{Hom}(N_1, N_2) & \xrightarrow{k_{2*}} & \mathrm{Hom}(N_1, X_2) & \xrightarrow{p_{2*}} & \mathrm{Hom}(N_1, M_2) & & \\
\downarrow \delta_1 & & \downarrow \delta_1 & & & & \\
\mathrm{Ext}^1(M_1, N_2) & \xrightarrow{k_{2*}} & \mathrm{Ext}^1(M_1, X_2) & & & &
\end{array}$$

Recall that $f \in \mathrm{Hom}(N_1, N_2)$ and $g \in \mathrm{Hom}(M_1, M_2)$. The assumption implies $\delta_1(f) = \delta_2(g)$. Hence $\delta_1 k_{2*}(f) = k_{2*} \delta_1(f) = k_{2*} \delta_2(g) = 0$. Hence there exists an element $\varphi' \in \mathrm{Hom}(X_1, X_2)$ such that $k_2^*(\varphi') = k_{2*}(f)$. Since $k_2^* p_{2*}(\varphi') = p_{2*} k_2^*(\varphi') = p_{2*} k_{2*}(f) = 0$, there exists $g' \in \mathrm{Hom}(M_1, M_2)$ such that $p_1^*(g') = p_{2*}(\varphi')$. Then $p_{2*}(\varphi') = p_1^*(g')$ and $p_1^*(\varphi') = k_{2*}(f)$. From the argument in the first part of this proof, we get $f_*a_1 = (g')^*a_2$, namely, we get $\delta_2(g') = \delta_1(f) = \delta_2(g)$. Hence there exists $r \in \mathrm{Hom}(M_1, X_2)$ such that $p_{2*}(r) = g - g'$. Set $\varphi = \varphi' + p_1^*(r)$. Then $p_{2*}(\varphi) = p_1^*(g)$ and $p_1^*(\varphi) = k_{2*}(f)$. This proves the lemma. $\square$

The following lemma is well-known. We give a proof for the same of completeness.

Lemma 3.2. *Let $x \in W$ and $s \in S$ such that $x < xs$.*

- (1) *We have $\dim \mathrm{Ext}^1(M(xs), M(x)) = 1$. The basis is given by the exact sequence $0 \rightarrow M(x) \rightarrow \theta_s(M(x)) \rightarrow M(xs) \rightarrow 0$.*
- (2) *The homomorphism $\mathrm{Ext}^1(M(xs), M(x)) \rightarrow \mathrm{Ext}^1(M(w_0), M(e))$ is injective and its image is independent of x .*

Proof. Let μ be the integral dominant element such that $\mathrm{Stab}_W(\mu) = \{e, s\}$. Then $\theta_s = T_\mu^\lambda T_\lambda^\mu$.

(1) We have

$$\mathrm{Ext}^i(M(xs), \theta_s(M(x))) = \mathrm{Ext}^i(T_\lambda^\mu(M(xs)), T_\lambda^\mu(M(x))) = \mathrm{Ext}^i(M(x\mu), M(x\mu))$$

It is one-dimensional if $i = 0$ and zero if $i > 0$. Hence from the exact sequence

$$0 \rightarrow M(x) \rightarrow \theta_s(M(x)) \rightarrow M(xs) \rightarrow 0,$$

we have an exact sequence

$$\begin{aligned}
0 \rightarrow \mathrm{Hom}(M(xs), M(x)) &\rightarrow \mathrm{Hom}(M(xs), \theta_s(M(x))) \\
&\rightarrow \mathrm{Hom}(M(xs), M(xs)) \rightarrow \mathrm{Ext}^1(M(xs), M(x)) \rightarrow 0.
\end{aligned}$$

By the dimension counting, $\mathrm{Hom}(M(xs), M(x)) \rightarrow \mathrm{Hom}(M(xs), \theta_s(M(x)))$ is isomorphic. Hence $\mathrm{Hom}(M(xs), M(xs)) \simeq \mathrm{Ext}^1(M(xs), M(x))$. The left hand side is one-dimensional.

- (2) First we prove that $\mathrm{Ext}^1(M(xs), M(x)) \rightarrow \mathrm{Ext}^1(M(w_0), M(e))$ is injective. The kernel of the homomorphism is $\mathrm{Hom}(M(xs), M(e)/M(x))$ by Lemma 2.1
- (3). We have an exact sequence $0 \rightarrow M(e) \rightarrow \theta_s(M(e)) \rightarrow M(s) \rightarrow 0$ and $0 \rightarrow M(x) \rightarrow \theta_s(M(x)) \rightarrow M(xs) \rightarrow 0$. Since $M(xs) \rightarrow M(s)$ is injective,

we have that $M(e)/M(x) \rightarrow \theta_s(M(e)/M(x))$ is injective by the snake lemma. Hence it is sufficient to prove that $\text{Hom}(M(xs), \theta_s(M(e)/M(x))) = 0$. We prove $\text{Hom}(M(x\mu), M(\mu)/M(x\mu)) = 0$. This follows from the following exact sequence

$$\begin{aligned} 0 \rightarrow \text{Hom}(M(x\mu), M(x\mu)) &\rightarrow \text{Hom}(M(x\mu), M(\mu)) \\ &\rightarrow \text{Hom}(M(x\mu), M(\mu)/M(x\mu)) \rightarrow \text{Ext}^1(M(x\mu), M(x\mu)) = 0. \end{aligned}$$

We prove the independence of x . We have the following diagram

$$\begin{array}{ccc} M(xs) & \longrightarrow & M(x) \\ \downarrow & & \downarrow \\ M(s) & \longrightarrow & M(e). \end{array}$$

Hence we get the following diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & M(x) & \longrightarrow & \theta_s(M(x)) & \longrightarrow & M(xs) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & M(e) & \longrightarrow & \theta_s(M(e)) & \longrightarrow & M(s) \longrightarrow 0. \end{array}$$

Therefore the image of

$$\text{Ext}^1(M(xs), M(x)) \rightarrow \text{Ext}^1(M(xs), M(e))$$

and

$$\text{Ext}^1(M(s), M(e)) \rightarrow \text{Ext}^1(M(xs), M(e))$$

coincide with each other by Lemma 3.1. Therefore, we get (2). $\square$

Take a basis v_s of $V(xs, x)$.

Lemma 3.3. *The set $\{v_s \mid s \in S\}$ is a basis of V .*

Proof. By Lemma 2.1 (1), we have the following commutative diagram

$$\begin{array}{ccc} \text{Ext}^1(M(s), M(e)) & \xrightarrow{\sim} & \bigoplus_{s' \in S} \text{Hom}(M(s), M(s')) \\ \downarrow & & \downarrow \\ \text{Ext}^1(M(w_0), M(e)) & \xrightarrow{\sim} & \bigoplus_{s' \in S} \text{Hom}(M(w_0), M(s')). \end{array}$$

We get the lemma. $\square$

We need the following lemma to define the action of $s \in S$ on V .

Lemma 3.4. *The homomorphism $\text{Ext}^1(M(w_0), M(e)) \rightarrow \text{Ext}^1(C_s(M(w_0)), M(e))$ is an isomorphism.*

Proof. From an exact sequence $0 \rightarrow M(w_0s)/M(w_0) \rightarrow C_s(M(w_0)) \rightarrow M(w_0) \rightarrow 0$, we get

$$\begin{aligned} \text{Hom}(M(w_0s)/M(w_0), M(e)) &\rightarrow \text{Ext}^1(M(w_0), M(e)) \\ &\rightarrow \text{Ext}^1(C_s(M(w_0)), M(e)) \rightarrow \text{Ext}^1(M(w_0s)/M(w_0), M(e)) \end{aligned}$$

It is sufficient to prove that $\text{Ext}^i(M(w_0s)/M(w_0), M(e)) = 0$ for $i = 0, 1$. We have the following exact sequence

$$\begin{aligned} 0 \rightarrow \text{Hom}(M(w_0s)/M(w_0), M(e)) &\rightarrow \text{Hom}(M(w_0s), M(e)) \\ &\rightarrow \text{Hom}(M(w_0), M(e)) \rightarrow \text{Ext}^1(M(w_0s)/M(w_0), M(e)) \\ &\rightarrow \text{Ext}^1(M(w_0s), M(e)) \rightarrow \text{Ext}^1(M(w_0), M(e)). \end{aligned}$$

The homomorphism $\text{Hom}(M(w_0s), M(e)) \rightarrow \text{Hom}(M(w_0), M(e))$ is an isomorphism. By Lemma 2.1 (3), the kernel of the homomorphism $\text{Ext}^1(M(w_0s), M(e)) \rightarrow \text{Ext}^1(M(w_0), M(e))$ is isomorphic to $\text{Hom}(M(w_0s), M(e)/M(e)) = 0$. We get the lemma. $\square$

Using this lemma, we consider the following homomorphism:

$$\begin{aligned} V = \text{Ext}^1(M(w_0), M(e)) &\rightarrow \text{Ext}^1(LC_s(M(w_0)), LC_s(M(e))) \\ &\simeq \text{Ext}^1(C_s(M(w_0)), M(s)) \\ &\rightarrow \text{Ext}^1(C_s(M(w_0)), M(e)) \\ &\simeq \text{Ext}^1(M(w_0), M(e)) = V. \end{aligned}$$

Define an action of s on V by the above homomorphism. In other words, for $v_i \in V$ and the corresponding exact sequences $0 \rightarrow M(e) \rightarrow X_i \rightarrow M(w_0) \rightarrow 0$, we have $s(v_1) = v_2$ if and only if there exists the following commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & C_s(M(e)) & \longrightarrow & C_s(X_1) & \longrightarrow & C_s(M(w_0)) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & M(e) & \longrightarrow & X_2 & \longrightarrow & M(w_0) \longrightarrow 0. \end{array}$$

Here, $C_s(M(e)) \rightarrow M(e)$ and $C_s(M(w_0)) \rightarrow M(w_0)$ are canonical homomorphisms.

The aim of this section is to show that this gives a structure of a W -module. Since $\{C_s\}$ satisfies the braid relations [MS05, Lemma 5.10], this action satisfies the braid relations.

Take an exact sequence $0 \rightarrow M(e) \rightarrow X \rightarrow M(w_0) \rightarrow 0$ and consider the following diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & M(e) & \longrightarrow & X & \longrightarrow & M(w_0) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \theta_s M(e) & \longrightarrow & \theta_s X & \longrightarrow & \theta_s M(w_0) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & M(e) & \longrightarrow & X & \longrightarrow & M(w_0) \longrightarrow 0, \end{array}$$

here the vertical maps are natural transformations. The compositions of left and right vertical morphisms are zero. Hence composition of the middle vertical morphisms factors through $X \rightarrow M(w_0)$ and $M(e) \rightarrow X$. By this way, we get an element of $\text{Hom}(M(w_0), M(e))$. This gives a morphism

$$\alpha_s: V = \text{Ext}^1(M(w_0), M(e)) \rightarrow \text{Hom}(M(w_0), M(e)).$$

Since we fix an inclusion $M(w_0) \hookrightarrow M(e)$, we regard $\alpha_s(v) \in K$ for $v \in V$. So we get $\alpha_s \in V^*$.

Lemma 3.5. *If $v \in \text{Ext}^1(M(w_0), M(e))$ satisfies $\alpha_s(v) = 0$, then $s(v) = v$.*

Proof. Let $0 \rightarrow M(e) \rightarrow X \rightarrow M(w_0) \rightarrow 0$ be the corresponding exact sequence. The assumption means that $X \rightarrow \theta_s(X) \rightarrow X$ is zero. Hence $\theta_s(X) \rightarrow X$ factors through $\theta_s(X) \rightarrow C_s(X)$. Namely, we get the following commutative diagram.

$$\begin{array}{ccccccc} 0 & \longrightarrow & C_s(M(e)) & \longrightarrow & C_s(X) & \longrightarrow & C_s(M(w_0)) \longrightarrow 0, \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & M(e) & \longrightarrow & X & \longrightarrow & M(w_0) \longrightarrow 0. \end{array}$$

This means that $s(v) = v$. $\square$

Lemma 3.6. *We have $\alpha_s(v_s) = 2$ and $s(v_s) = -v_s$.*

Proof. We consider v_s as an element of $V(w_0, w_0s)$. So we consider all things in $\{M \in \mathcal{O}_\lambda \mid [M : L(x)] = 0 \text{ for } x \neq w_0, w_0s\}$. This category is equivalent to the regular integral block of the BGG category of $\mathfrak{g} = \mathfrak{sl}_2(K)$. So we may assume that $\mathfrak{g} = \mathfrak{sl}_2(K)$.

Set $P_0 = M(e)$, $P_1 = \theta_s(M(e))$. Then $P_0 \oplus P_1$ is a projective generator of $\mathcal{O}_\lambda$. Set $A = \text{End}(P_0 \oplus P_1)$. Then $\mathcal{O}_\lambda$ is equivalent to the category of finitely generated right A -modules.

We have that $\dim \text{Hom}(P_0, P_0) = \dim \text{Hom}(P_0, P_1) = \dim \text{Hom}(P_1, P_0) = 1$, $\dim \text{Hom}(P_1, P_1) = 2$. Let $f: P_0 \rightarrow P_1$ and $g: P_1 \rightarrow P_0$ be the natural transformations. Then $\text{Hom}(P_0, P_1) = Kf$, $\text{Hom}(P_1, P_0) = Kg$ and $\text{Hom}(P_1, P_1) = \text{Kid} + Kfg$. Moreover, $gf = 0$. Set $e_i: P \rightarrow P_i$ be the projection. Then $A = Ke_0 + Ke_1 + Kf + Kg + Kfg$, here f stands for $P_0 \oplus P_1 \rightarrow P_0 \xrightarrow{f} P_1 \rightarrow P_0 \oplus P_1$. (g and fg are similar.)

First, we calculate $\alpha_s(v_s)$. To calculate it, we consider the following diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & P_0 & \xrightarrow{i} & P_1 & \xrightarrow{p} & P_1/P_0 \longrightarrow 0 \\ & & \downarrow a & & \downarrow b & & \downarrow c \\ 0 & \longrightarrow & P_1 & \xrightarrow{j} & P_1^{\oplus 2} & \xrightarrow{q} & P_1 \longrightarrow 0 \\ & & \downarrow s & & \downarrow t & & \downarrow u \\ 0 & \longrightarrow & P_0 & \xrightarrow{i} & P_1 & \xrightarrow{p} & P_1/P_0 \longrightarrow 0, \end{array}$$

here j is an inclusion $x \mapsto (x, 0)$ and q is second projection. Notice that we used $\theta_s(P_0) = \theta_s(P_1/P_0) = P_1$ and the second exact sequence splits.

We regard P_0 and P_1 are right A -modules. Then $P_0 = Ke_0 + Kg$, $P_1 = Ke_1 + Kf + Kfg$. We regard P_0 as a submodule of P_1 , namely, $P_0 = Kf + Kfg$. We fix a inclusion $M(w_0) = P_1/P_0 \rightarrow M(e) = P_0$ by $[e_1] \mapsto fg$, here $[e_1]$ is an image of e_1 . Set $h = fg$. Since $\text{End}(P_0) = \text{Kid} + Kh$, b is given by $b = (\alpha_1 + \beta_1h, \alpha_2 + \beta_2h)$ for some $\alpha_1, \alpha_2, \beta_1, \beta_2 \in K$. Since a and i are both natural transformation. So $a = i = f$. By $bi = ja$, we have $\alpha_1 = 1$. Since c is given by $[e_1] \mapsto fg$, we get $\alpha_2 = 0$ and $\beta_2 = 1$ by $cp = qb$. So we have $b = (1 + \beta_1h, h)$.

Next, we consider t . Define $\gamma_1, \gamma_2, \delta_1, \delta_2 \in K$ by $t = (\gamma_1 + \delta_1 h, \gamma_2 + \delta_2 h)$. The morphism s is given by $e_1 \mapsto fg$, $f \mapsto 0$ and $fg \mapsto 0$. Hence $tj = is$ implies that $\gamma_1 = 0$ and $\delta_1 = 1$. Since $uq = pt$, we get $\gamma_2 = 1$. Therefore, $t = (h, 1 + \delta_2 h)$. Hence the composition tb is given by $2h$. The image of e_1 under $P_1 \rightarrow \theta_s(P_1) \rightarrow P_1$ is $2h$. Since P_1/P_0 is given by $[e_1] \mapsto h$, this means $\alpha_s(v_s) = 2$.

Set $t' = (h, -1): \theta_s(P_1) \simeq P_1^{\oplus 2} \rightarrow P_1$. Then we have $t'b = 0$ and $(-u)q = pt'$. This means that there exists a following diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & C_s(P_0) & \longrightarrow & C_s(P_1) & \longrightarrow & C_s(P_1/P_0) \longrightarrow 0 \\ & & \downarrow \tilde{s} & & \downarrow \tilde{t}' & & \downarrow \widetilde{-u} \\ 0 & \longrightarrow & P_0 & \longrightarrow & P_1 & \longrightarrow & P_0/P_1 \longrightarrow 0. \end{array}$$

Here $\tilde{s}$ (resp. $\tilde{t}'$, $\widetilde{-u}$) is the morphisms induced by s (resp. t' , $-u$). Hence we have $s(v_s) = -v_s$. $\square$

Lemma 3.7. *For $v \in V$, we have $s(v) = v - \alpha_s(v)v_s$.*

Proof. Since $\alpha_s(v_s) \neq 0$, we have $V = Kv_s + \text{Ker } \alpha_s$. So we may assume that $v \in \text{Ker } \alpha_s$ or $v = v_s$. The first one is Lemma 3.5 and the second one is Lemma 3.6. $\square$

Proposition 3.8. *The action of $s \in S$ on V defines a representation of W .*

Proof. We should prove $s^2 = 1$. This follows from Lemma 3.7 and Lemma 3.6. $\square$

In fact, V is isomorphic to $\mathfrak{h}^*$ as a W -module. We prove it in the rest of this section.

Lemma 3.9. *If $xs > x$ and $ys > y$, then $V(xs, ys) = s(V(x, y))$.*

Proof. This follows from $C_s(M(x)) \simeq M(xs)$ and $C_s(M(y)) \simeq M(ys)$. $\square$

Lemma 3.10. *Let $m'_{s,s'}$ be an order of $ss' \in \text{GL}(V)$. Then there exists $v'_s \in K^\times v_s$ such that $s(v'_{s'}) = v'_{s'} - 2\cos(\pi/m'_{s,s'})v_s$ for all $s, s' \in S$.*

Proof. Since W is a finite group, V is defined over $\overline{\mathbb{Q}}$. Hence we may assume $K = \overline{\mathbb{Q}}$. Moreover, by the base change, we may assume that $K = \mathbb{C}$. Since W is a finite group, V has a W -invariant inner product $\langle \cdot, \cdot \rangle$. Take $v''_s \in \mathbb{C}v_s$ such that $\langle v''_s, v''_s \rangle = 2$. By Lemma 3.7, $s(v^*) = v^* - v^*(v_s)\alpha_s$ for $v^* \in V$. Hence the kernel of $s+1: V^* \rightarrow V^*$ is $K\alpha_s$. The linear map $v \mapsto \langle v, v''_s \rangle$ belongs to $\text{Ker}(s+1) = K\alpha_s$. Since $\langle v''_s, v''_s \rangle = \alpha_s(v_s)$, we have $s(v) = v - \langle v, v''_s \rangle v''_s$ by Lemma 3.7.

Put $a_{s,s'} = \langle v''_s, v''_{s'} \rangle$. Then we have $a_{s,s'} = \overline{a_{s',s}}$. By Lemma 3.7, $Kv'_s + Kv'_{s'}$ is stable under the action of ss' . Its characteristic polynomial is $t^2 + (2 - |a_{s,s'}|^2)t + 1$. Let α, α^{-1} be its eigenvalue. Since (W, S) is a Weyl group, $m'_{s,s'} \leq 6$. Hence $\alpha = e^{2\pi\sqrt{-1}/m'_{s,s'}}$ or $\alpha = e^{-2\pi\sqrt{-1}/m'_{s,s'}}$. We get $|a_{s,s'}|^2 = 2 + \alpha + \alpha^{-1} = 2 + 2\cos 2\pi/m'_{s,s'} = (2\cos(\pi/m'_{s,s'}))^2$. Hence we have $a_{s,s'} = 2e^{\sqrt{-1}\theta_{s,s'}} \cos(\pi/m'_{s,s'})$ for some $\theta_{s,s'} \in \mathbb{R}$. Recall that the Coxeter graph of (W, S) is a tree. Hence we can choose $v'_s \in e^{\sqrt{-1}\mathbb{R}}v''_s$ such that $a_{s,s'} = 2\cos(\pi/m'_{s,s'})$. So we get the lemma. $\square$

Lemma 3.11. *For $x, y \in W$ such that $xs > x \geq y < ys$, let A (resp. B) be the image of $\text{Ext}^1(M(xs), M(ys)) \rightarrow \text{Ext}^1(M(xs), M(y))$ (resp. $\text{Ext}^1(M(xs), M(x)) \rightarrow \text{Ext}^1(M(xs), M(y))$). Then $A \cap B = 0$ if and only if $x \not\geq ys$.*

Proof. By Lemma 3.2, B is one-dimensional. If $x \geq ys$, then we have a homomorphism $\text{Ext}^1(xs, x) \rightarrow \text{Ext}^1(xs, ys)$. So $B \subset A$.

On the other hand, assume that $A \cap B \neq 0$. Let $\mu \in \mathfrak{h}^*$ be a dominant integral element such that $\text{Stab}_W(\mu) = \{e, s\}$. Then T_λ^μ induces an homomorphism $\text{Ext}^1(M(xs), M(y)) \rightarrow \text{Ext}^1(M(xs\mu), M(y\mu))$. Since $T_\lambda^\mu(M(xs)) = T_\lambda^\mu(M(x)) = M(x\mu)$, the image of B under T_λ^μ is zero. Hence the homomorphism

$$\text{Ext}^1(M(xs), M(ys)) \rightarrow \text{Ext}^1(M(xs\mu), M(y\mu))$$

has a kernel. This homomorphism is equal to

$$\text{Ext}^1(M(xs), M(ys)) \rightarrow \text{Ext}^1(M(xs), \theta_s(M(y))).$$

From an exact sequence

$$0 \rightarrow M(ys) \rightarrow \theta_s(M(y)) \rightarrow C_s(M(ys)) \rightarrow 0,$$

we have $\text{Hom}(M(xs), C_s(M(ys))) \neq 0$. By the adjointness,

$$\text{Hom}(M(x), C_s(M(ys))) = \text{Hom}(K_s(M(xs)), M(ys)) = \text{Hom}(M(x), M(ys)).$$

Hence we have $x \geq ys$. $\square$

Lemma 3.12. *For $s, s' \in S$, let $\langle s, s' \rangle$ be the group generated by $\{s, s'\}$. Then for all $x, y \in w_0\langle s, s' \rangle$, $\text{Ext}^1(M(x), M(y)) \rightarrow \text{Ext}^1(M(w_0), M(e))$ is injective.*

Proof. Denote the irreducible quotient of $M(x)$ by $L(x)$. Let $w \in \langle s, s' \rangle$ be the longest element. Put

$$\mathcal{O}' = \{M \in \mathcal{O}_\lambda \mid [M : L(z)] = 0 \text{ for } z \notin w_0\langle s, s' \rangle\}.$$

Then $\mathcal{O}'$ is equivalent to the regular integral block of the BGG category of semisimple Lie algebra of rank 2. Applying Lemma 2.1 (3) to this category, the kernel of

$$\text{Ext}^1(M(x), M(y)) \rightarrow \text{Ext}^1(M(w_0), M(w_0w))$$

is isomorphic to $\text{Hom}(M(x), M(w_0w)/M(y))$. However, we have that $[M(w_0w) : L(x)] = [M(y) : L(x)] = 1$. Hence this space is zero. By Lemma 2.1 (3), the kernel of

$$\text{Ext}^1(M(w_0), M(w_0w)) \rightarrow \text{Ext}^1(M(w_0), M(e))$$

is isomorphic to $\text{Hom}(M(w_0), M(e)/M(w_0w))$. It is zero since $[M(e) : L(w_0)] = [M(w_0w) : L(w_0)] = 1$. $\square$

Remark 3.13. By Lemma 3.11 and Lemma 3.12, we have the following. For $x, y \in w_0\langle s, s' \rangle$ such that $xs > x \geq y < ys$, $v_s \in V(xs, ys)$ if and only if $x \geq ys$. Since $s(v_s) = -v_s$ and $V(xs, ys) = s(V(x, y))$ (Lemma 3.9), under the same conditions, we have $v_s \in V(x, y)$ if and only if $x \geq ys$.

Proposition 3.14. *The W -representation V is isomorphic to the geometric representation defined in [Bou02].*

Proof. Let $w' \in \langle s, s' \rangle$ be the longest element and put $w = w_0w'$. Let $m_{s, s'}$ be an order of $ss' \in W$. It is sufficient to prove that $m_{s, s'}$ is an order of $ss' \in \text{GL}(V)$.

Fix $s, s' \in S$ and set $m = m_{s, s'}$. Put $V' = Kv_s + Kv_{s'} = V(w_0, w)$. Then V' is $\langle s, s' \rangle$ -stable. Let n be an order of $ss' \in \text{GL}(V')$. Then $n|m$. We prove $n = m$. Since (W, S) is a Weyl group, $m = 2, 3, 4, 6$. If $m = 2, 3$, then there is nothing to prove.

Assume that $n = 2$ and $m \neq 2$. Then by Lemma 3.10, $s'v'_s = v'_s$. Therefore, $v_s \in s'V(ws, w) = V(wss', ws')$ by Lemma 3.9. By Remark 3.13, $ss' \geq s's$. This is a contradiction since $m \neq 2$.

Assume that $(n, m) = (3, 6)$. Then we have $s's(v'_{s'}) = -v'_s$. Hence, $v_s \in s'sV(ws', w) = V(ws'ss', wss')$. By Remark 3.13, $s'ss' \geq ss's$. This is a contradiction since $m = 6$. $\square$

This proposition says that $V \simeq \mathfrak{h}^*$ as W -module. Let α be a simple root. By the proof, v_{s_α} corresponds to α up to constant multiple.

4. REGULAR CASE

As in the previous section, fix a regular integral dominant element λ . We continue to use the notation in the previous section. In this section, we determine $V(x, y) \subset V$.

We need the graded BGG category. By Beilinson-Ginzburg-Soergel [BGS96], there exists a graded algebra $A = \bigoplus_{i \geq 0} A_i$ such that $\mathcal{O}_\lambda$ is equivalent to the category of A -modules. Let $\widetilde{\mathcal{O}}_\lambda$ be the category of graded A -modules. Then we have the forgetful functor $\widetilde{\mathcal{O}}_\lambda \rightarrow \mathcal{O}_\lambda$. For $M = \bigoplus_i M_i \in \widetilde{\mathcal{O}}_\lambda$, define $M\langle n \rangle \in \widetilde{\mathcal{O}}_\lambda$ by $(M\langle n \rangle)_i = M_{i-n}$. There exists a module $\widetilde{M}(x) \in \widetilde{\mathcal{O}}_\lambda$ such that its image in $\mathcal{O}_\lambda$ is isomorphic to $M(x)$ and it is unique up to grading shift [BGS96, 3.11]. Take a grading of $\widetilde{M}(x)$ such that its top is in degree 0. Set $\widetilde{M}^0(x) = \widetilde{M}(x)\langle \ell(x) \rangle$. Then by the proof of [Maz07, Corollary 23], if $x \geq y$, then there exists a degree zero injective homomorphism $\widetilde{M}^0(x) \rightarrow \widetilde{M}^0(y)$.

For $M, N \in \widetilde{\mathcal{O}}_\lambda$, let $\text{Hom}(M, N)_0$ be the space of homomorphisms of degree zero and $\text{Ext}^i(M, N)_0$ its derived functors. The following lemma is proved in [Maz07, Theorem 32]. However, the author think, although his proof is correct, the shift in the statement of the theorem is wrong. So we give a proof (it is the same as in the proof in [Maz07]).

Lemma 4.1. *We have $\text{Ext}^1(\widetilde{M}^0(x), \widetilde{M}^0(e)\langle i \rangle)_0 = 0$ if $i \neq 2$. Moreover, if $i = 2$, its dimension is equal to $\#\{s \in S \mid s \leq x\}$.*

Proof. We use the argument in the proof of Lemma 2.1 with a grading. Let $\widetilde{P}$ be the projective cover of $\widetilde{M}(w_0)$. Then it has a Verma flag M_i such that M_i/M_{i-1} is isomorphic to $\bigoplus_{\ell(x)=i} M(x)$ (here, we ignore the grading). The by Lemma 2.1 (1), we have $\text{Hom}(\widetilde{M}^0(x\lambda), M_1/M_0)_0 \simeq \text{Ext}^1(\widetilde{M}^0(x\lambda), M_0)$. We consider the grading of M_0 and M_1/M_0 . Take $k \in \mathbb{Z}$ such that $M_0 \simeq \widetilde{M}(e)\langle k \rangle$. Then the multiplicity of $\widetilde{M}(w_0)\langle k \rangle$ in $\widetilde{M}(e)$ is 1 by [BGS96, Theorem 3.11.4]. Since $\widetilde{M}(w_0)\langle \ell(w_0) \rangle \subset \widetilde{M}(e)$, we have $k = \ell(w_0)$. By the same argument, we have $M_1/M_0 \simeq \bigoplus_s \widetilde{M}(s)\langle \ell(w_0) - 1 \rangle$. So we have

$$\text{Ext}^1(\widetilde{M}^0(x), \widetilde{M}^0(e)\langle i \rangle) \simeq \text{Hom}\left(\widetilde{M}^0(x), \bigoplus_{s \in S} \widetilde{M}^0(s)\langle i - 2 \rangle\right).$$

We get the lemma. $\square$

We use the following abbreviations.

- $E^i(x\langle k \rangle, y\langle l \rangle) = \text{Ext}^i(\widetilde{M}^0(x)\langle k \rangle, \widetilde{M}^0(y)\langle l \rangle)_0$.
- $E^i(x\langle k \rangle, y/z\langle l \rangle) = \text{Ext}^i(\widetilde{M}^0(x)\langle k \rangle, \widetilde{M}^0(y)/\widetilde{M}^0(z)\langle l \rangle)_0$.

By the above lemma, $V(x, y)$ is the image of $E^1(x, y\langle 2 \rangle) \rightarrow E^1(w_0, e\langle 2 \rangle)$.

Lemma 4.2. *If $i \leq 1$, then $\text{Ext}(\widetilde{M}^0(x), \widetilde{M}^0(y)\langle i \rangle)_0 = 0$.*

Proof. We have $E^1(x, y\langle i \rangle) = E^0(x, e/y\langle i \rangle)$ by Lemma 2.1 (3) and the above lemma. Let L be the unique irreducible quotient of $\widetilde{M}^0(x)$. By [BGS96, Theorem 3.11.4], we have $[\widetilde{M}^0(e)/\widetilde{M}^0(y)\langle i \rangle : L] = 0$. Hence we have $E^0(x, e/y\langle i \rangle) = 0$. $\square$

There is a graded lift of θ_s [Str03]. We take a lift $\widetilde{\theta}_s$ of θ_s such that $\widetilde{\theta}_s$ is self-dual and there exist degree zero natural transformations $\text{Id}\langle 1 \rangle \rightarrow \widetilde{\theta}_s \rightarrow \text{Id}\langle -1 \rangle$. Let $\widetilde{C}_s$ (resp. $\widetilde{K}_s$) be the cokernel (resp. kernel) of $\text{Id}\langle 1 \rangle \rightarrow \widetilde{\theta}_s$ (resp. $\widetilde{\theta}_s \rightarrow \text{Id}\langle -1 \rangle$). Then $(\widetilde{C}_s, \widetilde{K}_s)$ is an adjoint pair. By [Str03, Theorem 3.6], we have the following formulas for $x \in W$ and $s \in S$ such that $xs > x$:

- $\widetilde{C}_s(\widetilde{M}^0(x)) = \widetilde{M}^0(xs)\langle -1 \rangle$ and $L^k \widetilde{C}_s(\widetilde{M}^0(x)) = 0$ for $k > 0$.
- We have an exact sequence $0 \rightarrow (\widetilde{M}^0(x)/\widetilde{M}^0(xs))\langle 1 \rangle \rightarrow \widetilde{C}_s(\widetilde{M}^0(xs)) \rightarrow \widetilde{M}^0(xs)\langle -1 \rangle \rightarrow 0$.
- $\widetilde{K}_s(\widetilde{M}^0(xs)) = \widetilde{M}^0(x)\langle 1 \rangle$ and $R^k \widetilde{K}_s(\widetilde{M}^0(xs)) = 0$ for $k > 0$.

We also have that $L\widetilde{C}_s$ gives an auto-equivalence of $D^b(\widetilde{\mathcal{O}}_\lambda)$ and its quasi-inverse functor is $R\widetilde{K}_s$.

Now we prove the main theorem in the regular case. We have already proved (1) (Lemma 3.9).

Theorem 4.3. *Let $x, y \in W$ and $s \in S$ such that $xs > x \geq y$.*

- (1) *If $ys < y$, then $V(xs, y) = s(V(x, ys))$.*
- (2) *If $ys > y$, then $V(xs, y) = Kv_s + s(V(x, y))$.*

Proof. (2) From (1), we have $s(V(x, y)) = V(xs, ys)$. Since $y < ys$, we have a homomorphism $\text{Ext}^1(M(xs), M(ys)) \rightarrow \text{Ext}^1(M(xs), M(y))$. Hence $V(xs, ys) \subset V(xs, y)$. We also have $\text{Ext}^1(M(xs), M(x)) \rightarrow \text{Ext}^1(M(xs), M(y))$. Since Kv_s is the image of $\text{Ext}^1(M(xs), M(x))$, the right hand side is contained in the left hand side.

From an exact sequence

$$0 \rightarrow (\widetilde{M}^0(y)/\widetilde{M}^0(ys))\langle 2 \rangle \rightarrow \widetilde{C}_s(\widetilde{M}^0(ys))\langle 1 \rangle \rightarrow \widetilde{M}^0(ys) \rightarrow 0,$$

we have an exact sequence

$$\begin{aligned} \text{Ext}^0(\widetilde{M}^0(xs), \widetilde{C}_s(\widetilde{M}^0(ys))\langle 1 \rangle)_0 &\rightarrow E^0(xs, ys) \\ &\rightarrow E^1(xs, y/ys\langle 2 \rangle) \rightarrow \text{Ext}^1(\widetilde{M}^0(xs), \widetilde{C}_s(\widetilde{M}^0(ys))\langle 1 \rangle)_0. \end{aligned}$$

Since $R\widetilde{K}_s$ is the quasi-inverse functor of $L\widetilde{C}_s$, for $i \geq 0$, we have

$$\begin{aligned} \text{Ext}^i(\widetilde{M}^0(xs), \widetilde{C}_s(\widetilde{M}^0(ys))\langle 1 \rangle)_0 &= \text{Hom}_{D^b(\widetilde{\mathcal{O}}_\lambda)}(\widetilde{M}^0(xs), L\widetilde{C}_s(\widetilde{M}^0(ys))\langle 1 \rangle[i])_0 \\ &= \text{Hom}_{D^b(\widetilde{\mathcal{O}}_\lambda)}(R\widetilde{K}_s(\widetilde{M}^0(xs)), \widetilde{M}^0(ys)\langle 1 \rangle[i])_0 \\ &= \text{Hom}_{D^b(\widetilde{\mathcal{O}}_\lambda)}(\widetilde{M}^0(x)\langle 1 \rangle, \widetilde{M}^0(ys)\langle 1 \rangle[i])_0 \\ &= \text{Ext}^i(\widetilde{M}^0(x), \widetilde{M}^0(ys))_0. \end{aligned}$$

If $i = 1$, then this is zero by Lemma 4.2. Hence we get an exact sequence

$$E^0(x, ys) \rightarrow E^0(xs, ys) \rightarrow E^1(xs, y/ys\langle 2 \rangle) \rightarrow 0.$$

Assume that $x \not\geq ys$. Then $E^0(x, ys) = 0$. Hence we have $\dim E^1(xs, y/ys\langle 2 \rangle) = \dim E^0(xs, ys) = 1$. From an exact sequence

$$E^1(xs, ys\langle 2 \rangle) \rightarrow E^1(xs, y\langle 2 \rangle) \rightarrow E^1(xs, y/ys\langle 2 \rangle),$$

the codimension of the image A of $E^1(xs, ys\langle 2 \rangle) \rightarrow E^1(xs, y\langle 2 \rangle)$ is less than or equal to 1. We also have that the image B of $E^1(xs, x\langle 2 \rangle) \rightarrow E^1(xs, ys\langle 2 \rangle)$ is one-dimensional by Lemma 3.2. By Lemma 3.11, $A \cap B = 0$. Hence $A + B = E^1(xs, y\langle 2 \rangle)$. This implies the theorem in this case.

Next assume that $x \geq ys$. Then $E^0(x, ys) \rightarrow E^0(xs, ys)$ is isomorphic. Hence $E^1(xs, y/ys\langle 2 \rangle) = 0$. Therefore, $E^1(xs, ys\langle 2 \rangle) \rightarrow E^1(xs, y\langle 2 \rangle)$ is surjective. This implies $V(xs, ys) \supset V(xs, y)$. $\square$

As a corollary, we can determine the dimension of $V(x, y)$. Let $R_{y,x}(q)$ be the polynomial defined in [KL79].

Theorem 4.4. *Assume that λ is regular. Then $\dim V_\lambda(x, y)$ is the coefficient of q in $(-1)^{\ell(y)-\ell(x)-1}R_{y,x}(q)$.*

Proof. Put $n_{y,x} = \dim V(x, y)$. Then by the above theorem and its proof, for $x, y \in W$, $s \in S$ such that $xs > x \geq y$, we have

- If $ys < y$, then $n_{xs,y} = n_{x,ys}$.
- If $ys > y$ and $x \geq ys$, then $n_{xs,y} = n_{x,y}$.
- If $ys > y$ and $x \not\geq ys$, then $n_{xs,y} \leq n_{x,y} + 1$.

Let $r_{y,x}$ be the coefficient of q in $(-1)^{\ell(y)-\ell(x)-1}R_{y,x}(q)$. By [KL79, (2.0.b), (2.0.c)], the constant term of $(-1)^{\ell(y)-\ell(x)-1}R_{y,x}(q)$ is 0 or 1 and it is 1 if and only if $y \leq x$. Hence using [KL79, (2.0.b), (2.0.c)], we have

- If $ys < y$, then $r_{xs,y} = r_{x,ys}$.
- If $ys > y$ and $x \geq ys$, then $r_{xs,y} = r_{x,y}$.
- If $ys > y$ and $x \not\geq ys$, then $r_{xs,y} = r_{x,y} + 1$.

Hence we get $n_{y,x} \leq r_{y,x}$. To prove $n_{y,x} = r_{y,x}$, it is sufficient to prove $n_{w_0,x} = r_{w_0,x}$. We prove this by backward induction on $\ell(x)$. By Lemma 4.1, we have $n_{w_0,x} = \#\{s' \in S \mid w_0s' \geq x\}$.

Take $s \in S$ such that $xs > x$. If $w_0s \geq xs$, then $r_{w_0,x} = r_{w_0s,x} = r_{w_0,xs} = n_{w_0,xs}$. If $w_0s \not\geq xs$, then $r_{w_0,x} = r_{w_0s,x} + 1 = r_{w_0,xs} + 1 = n_{w_0,xs} + 1$. We compare $X = \{s' \in S \mid w_0s' \geq x\}$ and $Y = \{s' \in S \mid w_0s' \geq xs\}$. Since $xs > x$, we have $Y \subset X$.

Assume $s' \in Y$ and $s' \neq s$. Then $s' \leq w_0xs$. Hence s' appears in a reduced expression of w_0xs . Since $s \neq s'$, s' appears in a reduced expression of w_0x . Hence $s' \in X$. This implies $X \cap (S \setminus \{s\}) = Y \cap (S \setminus \{s\})$.

Since $xs > x$, we have $w_0s \geq x$. Therefore, $s \in X$. Hence if $w_0s \geq xs$, we have $X = Y$, this implies $n_{w_0,x} = n_{w_0,xs}$. If $w_0s \not\geq xs$, we have $X = Y \amalg \{s\}$, this implies $n_{w_0,x} = n_{w_0,xs} + 1$. Therefore, $r_{w_0,x} = n_{w_0,x}$. $\square$

5. SINGULAR CASE

In this section, we fix a dominant integral (may be singular) element $\lambda \in \mathfrak{h}^*$. We also fix a regular integral dominant element $\lambda_0 \in \mathfrak{h}^*$. Then the translation functor $T_{\lambda_0}^\lambda$ is defined and it gives $V_{\lambda_0}(x, y) \rightarrow V_\lambda(x, y)$. Recall the notation $S_\lambda = \{s \in S \mid s(\lambda) = \lambda\}$.

In this section, we prove the following theorem.

Theorem 5.1. (1) *The homomorphism $V_{\lambda_0}(x, y) \rightarrow V_\lambda(x, y)$ induced by translation functor is surjective.*

(2) *The kernel of $V_{\lambda_0} \rightarrow V_\lambda$ is $\sum_{s \in S_\lambda} K v_s$.*

We use the notation $\widetilde{\mathcal{O}}_\lambda$, $M\langle n \rangle$ and $\text{Hom}(M, N)_0$ which we use in the previous section. Then using the same argument in [Str03], $T_{\lambda_0}^\lambda$ and $T_\lambda^{\lambda_0}$ have graded lifts $\widetilde{T}_{\lambda_0}^\lambda: \widetilde{\mathcal{O}}_{\lambda_0} \rightarrow \widetilde{\mathcal{O}}_\lambda$ and $\widetilde{T}_\lambda^{\lambda_0}: \widetilde{\mathcal{O}}_\lambda \rightarrow \widetilde{\mathcal{O}}_{\lambda_0}$, respectively.

Using the argument in [Str03], we can prove the following properties. Put $\widetilde{\theta} = \widetilde{T}_\lambda^{\lambda_0} \widetilde{T}_{\lambda_0}^\lambda$. Set $W_{S_\lambda} = \text{Stab}_W(\lambda)$ and let $w_\lambda \in W_{S_\lambda}$ be the longest element. Then we can take $\widetilde{T}_{\lambda_0}^\lambda$ and $\widetilde{T}_\lambda^{\lambda_0}$ such that $\widetilde{\theta}$ is self-dual and there exists a natural transformation $\text{Id}\langle \ell(w_\lambda) \rangle \rightarrow \widetilde{\theta}$ and $\widetilde{\theta} \rightarrow \text{Id}\langle -\ell(w_\lambda) \rangle$. Set $W(S_\lambda) = \{x \in W \mid xs > x \text{ for all } s \in S_\lambda\}$. Then for $x \in W(S_\lambda)$, $\widetilde{\theta}(\widetilde{M}^0(x\lambda_0))$ has a filtration M_i such that M_i/M_{i-1} is isomorphic to $\bigoplus_{\ell(w)=i, w \in W_{S_\lambda}} \widetilde{M}^0(xw\lambda_0)\langle \ell(w_\lambda) - 2\ell(w) \rangle$.

Proof of Theorem 5.1. (1) By Theorem 4.3, for $y \in W(S_\lambda)$ and $w \in W_{S_\lambda}$, we have $V(x, yw) \subset V(x, y) \subset V(x, yw) + \sum_{s \in S_\lambda} K v_s$. It is easy to see that v_s is in the kernel of $V_{\lambda_0} \rightarrow V_\lambda$. Hence we may assume that $y \in W(S_\lambda)$.

It is sufficient to prove that

$$\text{Ext}^1(\widetilde{M}^0(x\lambda_0), \widetilde{M}^0(y\lambda_0)\langle 2 \rangle)_0 \rightarrow \text{Ext}^1(\widetilde{M}^0(x\lambda_0), \widetilde{\theta}(\widetilde{M}^0(y\lambda_0))\langle 2 - \ell(w_\lambda) \rangle)_0$$

is surjective. Let M be the cokernel of

$$\widetilde{M}^0(y\lambda_0)\langle 2 \rangle \rightarrow \widetilde{\theta}(\widetilde{M}^0(y\lambda_0))\langle 2 - \ell(w_\lambda) \rangle.$$

Then it is sufficient to prove that $\text{Ext}^1(\widetilde{M}^0(x\lambda_0), M)_0 = 0$. As we mentioned above, M has a filtration $\{M'_i\}_{i \geq 1}$ such that $M'_i/M'_{i-1} \simeq \bigoplus_{\ell(w)=i, w \in W_\lambda} \widetilde{M}^0(yw\lambda_0)\langle 2 - 2\ell(w) \rangle$. By Lemma 4.2, we have $\text{Ext}^1(\widetilde{M}^0(x\lambda_0), \widetilde{M}^0(yw\lambda_0)\langle 2 - \ell(w) \rangle)_0 = 0$ if $\ell(w) > 0$. Hence we get (1).

(2) This follows from (1) and Lemma 2.1 (2). $\square$

REFERENCES

- [BGG71] I. N. Bernštein, I. M. Gel'fand, and S. I. Gel'fand, *Structure of representations that are generated by vectors of higher weight*, Funkcional. Anal. i Priložen. **5** (1971), no. 1, 1–9.
- [BGG76] I. N. Bernštein, I. M. Gel'fand, and S. I. Gel'fand, *A certain category of $\mathfrak{g}$ -modules*, Funkcional. Anal. i Priložen. **10** (1976), no. 2, 1–8.
- [BGS96] A. Beilinson, V. Ginzburg, and W. Soergel, *Koszul duality patterns in representation theory*, J. Amer. Math. Soc. **9** (1996), no. 2, 473–527.
- [Boe92] B. D. Boe, *A counterexample to the Gabber-Joseph conjecture*, Kazhdan-Lusztig theory and related topics (Chicago, IL, 1989), Contemp. Math., vol. 139, Amer. Math. Soc., Providence, RI, 1992, pp. 1–3.
- [Bou02] N. Bourbaki, *Lie groups and Lie algebras. Chapters 4–6*, Elements of Mathematics (Berlin), Springer-Verlag, Berlin, 2002, Translated from the 1968 French original by Andrew Pressley.
- [GJ81] O. Gabber and A. Joseph, *Towards the Kazhdan-Lusztig conjecture*, Ann. Sci. École Norm. Sup. (4) **14** (1981), no. 3, 261–302.
- [Jan79] J. C. Jantzen, *Moduln mit einem höchsten Gewicht*, Lecture Notes in Mathematics, vol. 750, Springer, Berlin, 1979.
- [KL79] D. Kazhdan and G. Lusztig, *Representations of Coxeter groups and Hecke algebras*, Invent. Math. **53** (1979), no. 2, 165–184.
- [Maz07] V. Mazorchuk, *Some homological properties of the category $\mathcal{O}$* , Pacific J. Math. **232** (2007), no. 2, 313–341.

- [MS05] V. Mazorchuk and C. Stroppel, *Translation and shuffling of projectively presentable modules and a categorification of a parabolic Hecke module*, Trans. Amer. Math. Soc. **357** (2005), no. 7, 2939–2973 (electronic).
- [Soe90] W. Soergel, *Kategorie $\mathcal{O}$, perverse Garben und Moduln über den Koinvarianten zur Weylgruppe*, J. Amer. Math. Soc. **3** (1990), no. 2, 421–445.
- [Str03] C. Stroppel, *Category $\mathcal{O}$: gradings and translation functors*, J. Algebra **268** (2003), no. 1, 301–326.
- [Ver68] D.-N. Verma, *Structure of certain induced representations of complex semisimple Lie algebras*, Bull. Amer. Math. Soc. **74** (1968), 160–166.